

007528
140693



Instructions for Use
SERNIVO™ (ser-ne-vo)
(betamethasone dipropionate)
Spray, 0.05%

Instructions for Use
SERNIVO™ (ser-ne-vo)
(betamethasone dipropionate)
Spray, 0.05%



007528
140693

Instructions for Use SERNIVO™ (ser-ne-vo) (betamethasone dipropionate) Spray, 0.05%

Important: SERNIVO Spray is for use on the skin only. Do not get SERNIVO Spray near or in your eyes, mouth, or vagina.

Read this “Instructions for Use” before you start using SERNIVO Spray and each time you get a refill. There may be new information. This information does not take the place of talking with your doctor about your medical condition or treatment.

Parts of the SERNIVO Spray bottle. (See Figure A)

Figure A



How to apply SERNIVO Spray:

Step 1: Shake the SERNIVO Spray bottle well. Remove the cap from the pump top.

Step 2: Hold the bottle in an upright position while pointing the opening of the pump top in the direction of the affected area. To spray, push down on the pump top. Apply SERNIVO Spray to the affected area as instructed by your doctor. **(See Figure B)**

Figure B



Step 3: Spray only enough SERNIVO Spray to cover the affected area, for example, the elbow **(See Figure C)**. Rub in SERNIVO Spray gently.

Figure C



Repeat Steps 2 and 3 to apply SERNIVO Spray to other affected areas as instructed by your doctor.

Step 4: After applying SERNIVO Spray, place the cap back onto the pump top. **(See Figure D)**

Figure D



How should I store SERNIVO Spray?

- Store SERNIVO Spray at room temperature between 68°F to 77°F (20°C to 25°C).
- Throw away (discard) any unused SERNIVO Spray after 28 days.

Keep SERNIVO Spray and all medicines out of the reach of children.

This “Instructions for Use” has been approved by the U.S. Food and Drug Administration.

Manufactured by:
DPT Laboratories, Ltd.,
San Antonio, TX 78215

Distributed by:
Promius Pharma, LLC.,
Princeton, NJ 08540

SERNIVO is a trademark of Promius Pharma, LLC.

Issued: 02/2016

007528
140693